

### Trend Study 17-47-02

Study site name: Tie Fork East.



Vegetation type: Mountain Brush.

Compass bearing: frequency baseline 0 degrees magnetic (line 2 @ 294°M, line 3 @ 325°M, line 4 @ 286°M).

Frequency belt placement: line 1 (11 & 95 ft), line 2 (59 ft), line 3 (34 ft), line 4 (71 ft). Rebar: belt 3 on 1 ft.

### LOCATION DESCRIPTION

From the intersection of Highway U.S. 6 and Tie Fork at Sky View in Spanish Fork Canyon, go north up to the railroad tracks. Cross the tracks and turn right. Follow the road along the railroad tracks for 0.65 miles. Stop at a pullout at the mouth of a small side canyon. Walk up the ridge to the west 200 yards to a witness post in a small rock outcrop on the bare ridgetop, by some mahogany. From the witness post, walk 18 paces north (3 degrees) to the 0-foot baseline stake.



Map Name: Tucker

Diagrammatic Sketch

Township 10S, Range 6E, Section 14

GPS: NAD 27, UTM 12S 4421995 N 482213 E

## DISCUSSION

### Tie Fork East - Trend Study No. 17-47

The Tie Fork range trend study (17-43) was established in the Tie Fork area in 1983. However, since the site was not considered representative of the large critical wintering area in Tie Fork, this study was established in 1989. The new study is on a more xeric site, supporting less oak and pinyon on a moderately steep slope. The area supports a scattered juniper community with a mountain brush understory. This site would be more representative of the winter range in the area. The lower end of Tie Fork is private land along the railroad. No livestock are currently grazed in the area but sheep are thought to trail through. The area receives considerably heavy winter use by deer, with light to moderate use by elk. Quadrat frequency of deer pellet groups was moderate in 1997 and 2002 at 38% and 33% respectively. A pellet group transect read on site in 2002 estimated 76 deer and 8 elk days use/acre (187 ddu/ha and 20 edu/ha). All of the deer and elk pellet groups appear to be from winter use.

This study is on a north facing slope at an elevation of 6,440 feet. The slope varies between 10% and 20%. Litter and organic matter buildup is good beneath shrub and tree crowns and combined with vegetative cover, helps to reduce erosion. However, there are still areas of localized erosion with active gullies forming below the site. The soil erosion condition class was determined to be slight in 2002. The soil is light gray in color and lacks structure. Soil textural analysis indicates a clay loam with a neutral reaction (pH 7.3). The effective rooting depth is 16 inches. Phosphorous is marginal at 8.3 ppm as values less than 10 ppm could be limiting to plant growth and development.

The browse community is a combination of large juniper and pinyon in association with an important shrub understory. The more prevalent juniper are largely unavailable due to height and past high-lining. Juniper trees average 25 to 30 feet in height with smaller trees in the understory. Density was estimated at 174 trees/acre in 2002 from point-center quarter data. Diameter of juniper averaged nearly 8 inches. Pinyon are less common on this site with a density of 30 trees/acre estimated in 2002. Juniper and pinyon canopy cover is variable but averaged almost 5% for juniper and nearly 2% for pinyon over the whole site.

Several preferred forage species occur in the understory including serviceberry, mountain big sagebrush, true mountain mahogany, and bitterbrush. Saskatoon serviceberry density was estimated about 300 plants/acre in 1997 and 2002. It displayed heavy use in 2002, although vigor was normal. True mountain mahogany is the most abundant palatable species. Density was estimated at 700 plants/acre in 2002. Utilization has been consistently heavy since 1989, yet most plants have good vigor. The number of decadent plants has ranged from 11% in 1997 to 23% in 2002.

Sagebrush on the site has some characteristics of basin big sagebrush (*Artemisia tridentata tridentata*) but was classified as Mountain big sagebrush (*Artemisia tridentata vaseyana*) in 2002. It had a low density estimated at only 260 plants/acre in 2002. Utilization was moderate to heavy in 1997 and 2002 and the number of decadent plants was moderately high at 50% and 38% respectively. Bitterbrush are highly preferred but only a few heavily utilized plants occur on the site.

Snowberry is abundant but less preferred. It had a density estimated at 5,300 plants/acre with mostly light utilization in 1997. Density increased to 6,200 plants/acre in 2002, and use continued to be mostly light. A few oak clones occur on site. These average about three feet in height with light to moderate use on available plants. Other scattered browse species include white rubber rabbitbrush, stickyleaf rabbitbrush, Wood's rose, and gray horsebrush.

Total herbaceous cover is relatively low on this site with grasses and forbs providing only 9% cover in 2002. The most abundant grasses are Indian ricegrass, a *Carex*, and bluebunch wheatgrass. Indian ricegrass and *Carex* remained at stable frequencies in 2002, but bluebunch wheatgrass significantly declined. Cheatgrass is present but not in high amounts. Forb cover is sparse with high diversity but low abundance. Several annual species were encountered, with many of the species being increasers or invaders including musk thistle.

## 1989 APPARENT TREND ASSESSMENT

With continued soil loss and no signs of an increasing understory or herbaceous component, the soil trend appears to be declining. The vegetative trend also has some downward indicators such as increasing juniper and continued heavy use of the preferred browse. Still, there is fair reproduction and significant amounts of unutilized browse forage (especially on the sagebrush).

## 1997 TREND ASSESSMENT

Erosion is still apparent, but is not excessive. This vegetative type will likely always have some erosion occurring, so establishment of herbaceous understory should be encouraged. Soil trend is up slightly due to a decline in bare ground and an increase in the sum of nested frequency for grasses and forbs. The browse trend is stable. There is little change in any of the browse species. The mountain big sagebrush population is 50% decadent with similar utilization and vigor as reported in 1989. It is still only a minor component of the browse composition at this time. True mountain mahogany has remained stable with moderate to heavy use, good vigor, and low decadence. The herbaceous understory trend is slightly upward with an increase in nested frequency for both perennial grasses and forbs.

### TREND ASSESSMENT

soil - up slightly (4)

browse - stable (3)

herbaceous understory - slightly upward (4)

## 2002 TREND ASSESSMENT

Trend for soil is down slightly. Cover of bare ground has increased from 15% to 25%, while litter cover has declined slightly. In addition, total herbaceous cover has declined from 14% to 9% and the sum of nested frequency for perennial grasses and forbs has declined. Current erosion is occurring but it is not severe and the erosion condition class was determined to be slight. Trend for browse is stable but showing the effects of drought. Serviceberry, mountain big sagebrush, and true mountain mahogany are the key browse species. Densities have remained relatively stable while use is more heavy. Sagebrush and mahogany display increased poor vigor and the number of decadent mahogany has increased from 11% to 23%. However, recruitment appears adequate to maintain their respective populations. Annual leader growth for mahogany was estimated at only 1.6 inches. Most plants showed little growth and many terminal buds were shriveling up by July 3<sup>rd</sup> due to the dry conditions. Trend for the herbaceous understory is down slightly. Sum of nested frequency for perennial grasses and forbs has declined and frequency of bluebunch wheatgrass declined significantly. The other abundant perennial grass, Indian ricegrass, has remained stable. Nested frequency of the most abundant forb, Wasatch penstemon, also declined significantly. Another negative indicator with respect to the herbaceous understory is the continued increase in cover of juniper.

### TREND ASSESSMENT

soil - down slightly (2)

browse - stable (3)

herbaceous understory - down slightly (2)

HERBACEOUS TRENDS --  
Herd unit 17 , Study no: 47

T y p e	Species	Nested Frequency			Quadrat Frequency			Average Cover %	
		'89	'97	'02	'89	'97	'02	'97	'02
G	Agropyron spicatum	<sub>a</sub> 8	<sub>b</sub> 42	<sub>a</sub> 12	3	20	5	2.20	.31
G	Bromus tectorum (a)	-	<sub>b</sub> 101	<sub>a</sub> 11	-	38	5	.48	.03
G	Carex spp.	<sub>a</sub> 6	<sub>ab</sub> 25	<sub>b</sub> 34	4	11	13	1.14	.78
G	Oryzopsis hymenoides	<sub>b</sub> 121	<sub>a</sub> 77	<sub>a</sub> 77	49	30	33	3.03	3.55
G	Poa fendleriana	-	-	4	-	-	2	-	.16
G	Poa pratensis	24	19	10	8	7	5	.13	.10
G	Poa secunda	-	6	-	-	2	-	.30	-
G	Sitanion hystrix	7	16	11	4	7	6	.13	.14
G	Stipa columbiana	10	7	8	3	3	3	.41	.04
G	Stipa lettermani	<sub>a</sub> 1	<sub>b</sub> 20	<sub>ab</sub> 8	1	8	3	.67	.21
Total for Annual Grasses		0	101	11	0	38	5	0.48	0.03
Total for Perennial Grasses		177	212	164	72	88	70	8.05	5.30
Total for Grasses		177	313	175	72	126	75	8.53	5.33
F	Achillea millefolium	5	6	4	2	2	2	.18	.01
F	Agoseris glauca	-	1	11	-	1	4	.00	.02
F	Antennaria rosea	7	7	-	2	2	-	.41	-
F	Astragalus convallarius	3	8	10	2	5	5	.02	.08
F	Castilleja linariaefolia	2	-	4	2	-	2	.01	.03
F	Carduus nutans (a)	<sub>a</sub> -	<sub>b</sub> 13	<sub>b</sub> 23	-	7	11	.42	.30
F	Chenopodium album (a)	-	1	-	-	1	-	.00	-
F	Chaenactis douglasii	7	4	-	4	2	-	.03	-
F	Cirsium spp.	4	21	2	4	11	1	.28	.03
F	Collinsia parviflora (a)	-	13	8	-	6	3	.03	.01
F	Cryptantha spp.	4	1	12	2	1	6	.03	.22
F	Cynoglossum officinale	<sub>c</sub> 107	<sub>b</sub> 50	<sub>a</sub> 2	43	19	1	.99	.00
F	Delphinium nuttallianum	-	1	-	-	1	-	.00	-
F	Descurainia pinnata (a)	<sub>a</sub> -	<sub>b</sub> 29	<sub>a</sub> 2	-	13	1	.09	.00
F	Epilobium brachycarpum (a)	-	3	4	-	1	2	.00	.03
F	Erigeron spp.	-	2	-	-	2	-	.01	-
F	Gayophytum ramosissimum (a)	-	-	1	-	-	1	-	.03
F	Hackelia patens	<sub>a</sub> -	<sub>b</sub> 16	<sub>b</sub> 21	-	7	11	.41	.44
F	Lappula occidentalis (a)	-	1	-	-	1	-	.00	-
F	Lactuca serriola	-	-	2	-	-	1	-	.00
F	Lithospermum ruderales	-	-	2	-	-	1	-	.06
F	Machaeranthera canescens	11	27	9	6	12	6	.13	.03
F	Melilotus officinalis	-	1	-	-	1	-	.00	-
F	Penstemon cyananthus	<sub>a</sub> 58	<sub>b</sub> 101	<sub>a</sub> 69	29	43	31	1.96	2.35

T y p e	Species	Nested Frequency			Quadrat Frequency			Average Cover %	
		'89	'97	'02	'89	'97	'02	'97	'02
F	Penstemon humilis	<sub>b</sub> 16	<sub>a</sub> -	<sub>a</sub> 2	6	-	1	-	.03
F	Phlox hoodii	-	-	1	-	-	1	-	.00
F	Phlox longifolia	3	3	-	1	2	-	.01	-
F	Ranunculus testiculatus (a)	-	3	3	-	1	1	.00	.00
F	Schoenocrambe linifolia	<sub>a</sub> -	<sub>b</sub> 16	<sub>ab</sub> 9	-	9	4	.16	.04
F	Senecio multilobatus	3	2	-	1	1	-	.03	-
F	Streptanthus cordatus	-	-	-	-	-	-	.00	-
F	Taraxacum officinale	-	2	-	-	1	-	.00	-
F	Tragopogon dubius	-	6	3	-	4	2	.04	.01
F	Unknown forb-perennial	3	-	-	1	-	-	-	-
F	Verbascum thapsus	-	7	4	-	3	2	.07	.18
Total for Annual Forbs		0	63	41	0	30	19	0.56	0.39
Total for Perennial Forbs		233	282	167	105	129	81	4.82	3.57
Total for Forbs		233	345	208	105	159	100	5.39	3.97

Values with different subscript letters are significantly different at alpha = 0.10

#### BROWSE TRENDS --

Herd unit 17 , Study no: 47

T y p e	Species	Strip Frequency		Average Cover %	
		'97	'02	'97	'02
B	Amelanchier alnifolia	13	9	.21	.09
B	Artemisia tridentata vaseyana	8	8	.45	.36
B	Cercocarpus montanus	22	26	2.58	2.12
B	Chrysothamnus nauseosus albicaulis	1	2	-	.00
B	Chrysothamnus viscidiflorus viscidiflorus	4	8	.06	.06
B	Juniperus osteosperma	10	12	2.23	4.59
B	Mahonia repens	1	1	-	-
B	Opuntia spp.	3	1	-	-
B	Pinus edulis	0	2	-	-
B	Purshia tridentata	1	0	.01	-
B	Quercus gambelii	9	8	2.01	1.16
B	Ribes spp.	0	1	-	-
B	Rosa woodsii	11	12	.57	.22
B	Symphoricarpos oreophilus	58	67	11.46	13.06
B	Tetradymia canescens	2	3	-	.15
Total for Browse		143	160	19.61	21.84

CANOPY COVER -- LINE INTERCEPT

Herd unit 17 , Study no: 47

Species	Percent Cover	
	'97	'02
Amelanchier utahensis	-	.08
Artemisia tridentata vaseyana	-	.25
Cercocarpus montanus	-	2.67
Chrysothamnus viscidiflorus viscidiflorus	-	.42
Juniperus osteosperma	5.8	4.42
Pinus edulis	2.0	1.58
Quercus gambelii	4.4	2.00
Rosa woodsii	-	.07
Symphoricarpos oreophilus	-	18.33

Key Browse Annual Leader Growth

Herd unit 17 , Study no: 47

Species	Average leader growth (in) '02
Cercocarpus montanus	1.6

Point-Quarter Tree Data

Herd unit 17, Study no: 47

Species	Trees per Acre		Average diameter (in)	
	'97	'02	'97	'02
Juniperus osteosperma	106	174	8.9	7.8
Pinus edulis	22	30	8.8	4.8

BASIC COVER --

Herd unit 17 , Study no: 47


Cover Type	Nested Frequency		Average Cover %		
	'97	'02	'89	'97	'02
Vegetation	314	244	7.25	31.06	31.76
Rock	115	155	2.25	4.31	4.36
Pavement	176	187	13.50	4.74	8.36
Litter	395	379	50.50	50.47	48.48
Cryptogams	39	58	0	.70	3.32
Bare Ground	203	277	26.50	15.30	24.68

# SOIL ANALYSIS DATA --

Herd Unit 17, Study no: 47, Tie Fork East

Effective rooting depth (in)	Temp °F (depth)	pH	%sand	%silt	%clay	%OM	PPM P	PPM K	dS/m
16.1	45.4 (15.4)	7.3	26.7	34.4	38.8	4.5	8.3	112.0	.5

## Stoniness Index



## PELLET GROUP FREQUENCY --

Herd unit 17 , Study no: 47

Type	Quadrat Frequency	
	'97	'02
Sheep	-	1
Rabbit	23	21
Elk	12	3
Deer	38	33

Pellet Transect	
Pellet Groups per Acre	Days Use per Acre (ha)
'02	'02
-	-
-	-
104	8 (20)
983	76 (187)

## BROWSE CHARACTERISTICS --

Herd unit 17 , Study no: 47

A G R E	Y R E	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Amelanchier alnifolia																		
Y	89	2	-	-	4	-	-	1	-	-	7	-	-	-	233		7	
	97	9	-	-	4	-	-	-	-	-	13	-	-	-	260		13	
	02	-	-	1	5	-	-	-	-	-	6	-	-	-	120		6	
M	89	-	-	1	2	-	-	-	-	-	3	-	-	-	100	27	20	
	97	-	-	-	-	1	1	-	-	-	2	-	-	-	40	26	29	
	02	-	-	3	4	1	-	-	-	-	8	-	-	-	160	22	18	
D	89	-	1	-	-	2	-	-	-	-	2	-	-	1	100		3	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	20		1	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		23%			08%			08%			-31%							
'97		07%			07%			00%			- 7%							
'02		07%			29%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	433	Dec:	23%			
												'97	300		0%			
												'02	280		0%			
Artemisia tridentata vaseyana																		
S	89	1	-	-	1	-	-	-	-	-	2	-	-	-	66		2	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
Y	89	4	-	-	1	-	-	-	-	-	5	-	-	-	166		5	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	2	-	-	-	-	-	-	-	-	2	-	-	-	40		2	
M	89	3	-	-	-	-	-	-	-	-	3	-	-	-	100	20	10	
	97	1	2	1	-	-	-	-	-	-	4	-	-	-	80	27	32	
	02	4	1	1	-	-	-	-	-	-	6	-	-	-	120	23	31	
D	89	5	1	1	-	-	-	-	-	-	6	-	-	1	233		7	
	97	2	1	-	1	-	-	-	-	-	3	-	-	1	80		4	
	02	-	3	2	-	-	-	-	-	-	2	-	-	3	100		5	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	260		13	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	320		16	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		07%			07%			07%			-68%							
'97		38%			13%			13%			+38%							
'02		31%			23%			23%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	499	Dec:	47%			
												'97	160		50%			
												'02	260		38%			



A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Cercocarpus montanus																		
S	89	1	-	-	-	-	-	-	-	-	1	-	-	-	33		1	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
Y	89	6	-	-	-	-	-	-	-	-	6	-	-	-	200		6	
	97	2	-	-	-	-	1	-	-	-	3	-	-	-	60		3	
	02	-	-	2	-	-	-	-	-	-	2	-	-	-	40		2	
M	89	-	-	10	-	-	-	-	-	-	10	-	-	-	333	67	79	
	97	1	2	4	1	7	6	-	-	-	21	-	-	-	420	33	29	
	02	6	1	16	-	2	-	-	-	-	24	-	1	-	500	31	26	
D	89	-	1	2	-	-	-	-	-	-	2	-	1	-	100		3	
	97	-	-	-	1	1	1	-	-	-	3	-	-	-	60		3	
	02	-	-	6	-	-	2	-	-	-	5	-	1	2	160		8	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	20		1	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	20		1	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		05%			63%			05%			-15%							
'97		37%			44%			00%			+23%							
'02		09%			74%			11%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	633	Dec:	16%			
												'97	540		11%			
												'02	700		23%			
Chrysothamnus nauseosus albicaulis																		
Y	89	1	-	-	-	-	-	-	-	-	1	-	-	-	33		1	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20		1	
M	89	1	-	1	-	-	-	-	-	-	2	-	-	-	66	35	22	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0	24	23	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0	24	23	
D	89	1	1	1	-	-	-	-	-	-	3	-	-	-	100		3	
	97	1	-	-	-	-	-	-	-	-	1	-	-	-	20		1	
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20		1	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		17%			33%			00%			-90%							
'97		00%			00%			00%			+50%							
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	199	Dec:	50%			
												'97	20		100%			
												'02	40		50%			

A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Chrysothamnus viscidiflorus viscidiflorus																		
Y	89	27	-	-	4	-	-	-	-	-	31	-	-	-	1033		31	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20		1	
M	89	20	-	-	-	-	-	1	-	-	21	-	-	-	700	18	24	
	97	5	-	-	-	-	-	-	-	-	5	-	-	-	100	12	12	
	02	9	4	-	-	-	-	-	-	-	10	3	-	-	260	11	16	
D	89	7	-	-	-	-	-	-	-	-	7	-	-	-	233		7	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%			-95%							
'97		00%			00%			00%			+64%							
'02		29%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	1966	Dec:	12%			
												'97	100		0%			
												'02	280		0%			
Juniperus osteosperma																		
S	89	-	-	-	1	-	-	-	-	-	1	-	-	-	33		1	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	2	-	-	-	-	-	-	-	-	2	-	-	-	40		2	
Y	89	1	-	-	1	-	-	-	-	-	2	-	-	-	66		2	
	97	6	-	-	-	-	-	-	-	-	6	-	-	-	120		6	
	02	4	-	-	-	-	-	-	-	-	3	-	1	-	80		4	
M	89	-	-	-	-	-	-	-	-	1	1	-	-	-	33	197	122	
	97	4	-	-	-	-	-	-	1	-	5	-	-	-	100	3	5	
	02	6	-	-	-	-	-	-	1	-	6	-	1	-	160	-	-	
D	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	2	-	-	-	-	-	-	-	-	-	-	1	1	40		2	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			33%			00%			+55%							
'97		00%			00%			00%			+21%							
'02		00%			00%			29%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	99	Dec:	0%			
												'97	220		0%			
												'02	280		14%			
Mahonia repens																		
M	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	
	97	1	-	-	-	-	-	-	-	-	1	-	-	-	20	3	5	
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20	2	3	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%										
'97		00%			00%			00%			+ 0%							
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	0	Dec:	-			
												'97	20		-			
												'02	20		-			

A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Opuntia spp.																		
Y	89	1	-	-	-	-	-	-	-	-	1	-	-	-	33		1	
	97	1	-	-	-	-	-	-	-	-	1	-	-	-	20		1	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
M	89	2	-	-	-	-	-	-	-	-	2	-	-	-	66	5	6	
	97	1	-	-	-	-	-	-	-	-	1	-	-	-	20	5	7	
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20	6	16	
D	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	1	-	-	-	-	-	-	-	-	-	-	-	1	20		1	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	20		1	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%			-39%							
'97		00%			00%			33%			-67%							
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	99	Dec:	0%			
												'97	60		33%			
												'02	20		0%			
Pinus edulis																		
Y	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	1	-	-	1	-	-	-	-	-	2	-	-	-	40		2	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%										
'97		00%			00%			00%										
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	0	Dec:	-			
												'97	0		-			
												'02	40		-			
Purshia tridentata																		
M	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	
	97	-	1	-	-	-	-	-	-	-	1	-	-	-	20	12	24	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0	13	33	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	20		1	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%										
'97		100%			00%			00%										
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	0	Dec:	-			
												'97	20		-			
												'02	0		-			

A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Quercus gambelii																		
S	89	1	-	-	-	-	-	-	-	-	1	-	-	-	33			1
	97	2	-	-	-	-	-	-	-	-	2	-	-	-	40			2
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
Y	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
	97	7	6	-	1	-	-	-	-	-	14	-	-	-	280			14
	02	4	-	-	2	-	-	-	-	-	6	-	-	-	120			6
M	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	0
	97	20	3	-	-	-	-	3	-	-	26	-	-	-	520	49	29	26
	02	23	12	-	7	-	-	-	-	-	35	-	7	-	840	33	18	42
D	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
	97	-	1	-	-	-	-	-	-	-	1	-	-	-	20			1
	02	2	-	-	-	-	-	-	-	-	-	-	1	1	40			2
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	60			3
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	160			8
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%										
'97		24%			00%			00%			+18%							
'02		24%			00%			18%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	0	Dec:	0%			
												'97	820		2%			
												'02	1000		4%			
Ribes spp.																		
Y	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0			0
	02	1	-	-	-	-	-	-	-	-	1	-	-	-	20			1
M	89	-	-	-	-	2	-	-	-	-	2	-	-	-	66	18	22	2
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	0
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	0
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		100%			00%			00%										
'97		00%			00%			00%										
'02		00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	66	Dec:	-			
												'97	0		-			
												'02	20		-			

A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Rosa woodsii																		
S	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	4	-	-	-	-	-	-	-	-	-	-	-	-	80		4	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
Y	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	12	-	-	11	-	-	-	-	-	23	-	-	-	460		23	
	02	7	-	-	7	-	-	-	-	-	14	-	-	-	280		14	
M	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	0	
	97	4	-	-	5	-	-	-	-	-	9	-	-	-	180	22	9	
	02	6	-	-	-	-	-	-	-	-	6	-	-	-	120	6	6	
D	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	1	-	-	-	1	-	-	-	20		1	
	02	1	-	-	-	-	-	-	-	-	-	-	-	1	20		1	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		00%			00%			00%										
'97		00%			03%			00%			-36%							
'02		00%			00%			05%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	0	Dec:	0%			
												'97	660		3%			
												'02	420		5%			
Symphoricarpos oreophilus																		
S	89	1	-	-	-	-	-	-	-	-	1	-	-	-	33		1	
	97	3	4	-	-	-	-	-	-	-	6	-	-	1	140		7	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
Y	89	8	2	1	-	-	-	12	-	-	23	-	-	-	766		23	
	97	29	-	-	-	-	-	-	-	-	29	-	-	-	580		29	
	02	12	-	-	-	-	-	-	-	-	12	-	-	-	240		12	
M	89	36	21	5	33	6	-	15	-	-	116	-	-	-	3866	20	116	
	97	148	40	11	37	-	-	-	-	-	236	-	-	-	4720	43	236	
	02	240	17	25	-	-	-	-	-	-	267	15	-	-	5640	18	282	
D	89	-	2	1	-	-	-	-	-	-	3	-	-	-	100		3	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	02	14	-	2	-	-	-	-	-	-	11	-	-	5	320		16	
X	89	-	-	-	-	-	-	-	-	-	-	-	-	-	0		0	
	97	-	-	-	-	-	-	-	-	-	-	-	-	-	80		4	
	02	-	-	-	-	-	-	-	-	-	-	-	-	-	40		2	
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
'89		22%			05%			00%			+11%							
'97		15%			04%			00%			+15%							
'02		05%			09%			02%										
Total Plants/Acre (excluding Dead & Seedlings)												'89	4732	Dec:	2%			
												'97	5300		0%			
												'02	6200		5%			

A G E	Y R	Form Class (No. of Plants)									Vigor Class				Plants Per Acre	Average (inches) Ht. Cr.		Total
		1	2	3	4	5	6	7	8	9	1	2	3	4				
Tetradymia canescens																		
M	'89	-	-	-	-	-	-	-	-	-	-	-	-	-	0	-	-	0
	'97	-	3	-	-	-	-	-	-	-	-	3	-	-	60	13	11	3
	'02	3	-	-	-	-	-	-	-	-	2	1	-	-	60	13	16	3
% Plants Showing		<u>Moderate Use</u>			<u>Heavy Use</u>			<u>Poor Vigor</u>			<u>%Change</u>							
		'89 00%			00%			00%										
		'97 100%			00%			00%			+ 0%							
		'02 00%			00%			00%										
Total Plants/Acre (excluding Dead & Seedlings)												'89		0	Dec:	-		
												'97		60		-		
												'02		60		-		